

Research article

Development and application of KASP marker for high throughput detection of *AhFAD2* mutation in peanut

Shuzhen Zhao, Aiqin Li, Changsheng Li, Han Xia, Chuanzhi Zhao, Ye Zhang, Lei Hou, Xingjun Wang*

Biotechnology Research Center, Shandong Academy of Agricultural Sciences, Shandong Provincial Key Laboratory of Crop Genetic Improvement, Ecology and Physiology, Jinan 250100, PR China

ARTICLE INFO

Article history:

Received 27 June 2016

Accepted 24 October 2016

Available online 16 November 2016

Keywords:

Arachis hypogaea

Fatty acid desaturase

High oleic acid

Competitive allele specific PCR

Linoleic acid

Marker assisted selection

Molecular breeding

Mutation in peanut

Nucleotide substitution

Oilseed crop

Peanut oil

ABSTRACT

Background: Cultivated peanut (*Arachis hypogaea* L.) is a major oilseed crop worldwide. Fatty acid composition of peanut oil may affect the flavor and shelf life of the resulting food products. Oleic acid and linoleic acid are the major fatty acids of peanut oil. The conversion from oleic acid to linoleic acid is controlled by the $\Delta 12$ fatty acid desaturase (FAD) encoded by *AhFAD2A* and *AhFAD2B*, two homoeologous genes from A and B subgenomes, respectively. One nucleotide substitution (G:C \rightarrow A:T) of *AhFAD2A* and an "A" insertion of *AhFAD2B* resulted in high-oleic acid phenotype. Detection of *AhFAD2* mutation had been achieved by cleaved amplified polymorphic sequence (CAPS), real-time polymerase chain reaction (qRT-PCR) and allele-specific PCR (AS-PCR). However, a low cost, high throughput and high specific method is still required to detect *AhFAD2* genotype of large number of seeds. Kompetitive allele specific PCR (KASP) can detect both alleles in a single reaction. The aim of this work is to develop KASP for detection *AhFAD2* genotype of large number of breeding materials.

Results: Here, we developed a KASP method to detect the genotypes of progenies between high oleic acid peanut and common peanut. Validation was carried out by CAPS analysis. The results from KASP assay and CAPS analysis were consistent. The genotype of 18 out of 179 BC₄F₂ seeds was aabb.

Conclusions: Due to high accuracy, time saving, high throughput feature and low cost, KASP is more suitable for determining *AhFAD2* genotype than other methods.

© 2016 Pontificia Universidad Católica de Valparaíso. Production and hosting by Elsevier B.V. All rights reserved. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Cultivated peanut (*Arachis hypogaea* L.) is one of the most important oil crops worldwide. Oleic acid and linoleic acid are the major fatty acids of peanut oil, accounting for approximately 80% of the oil composition [1]. Linoleic acid contains two double bonds at the $\Delta 9$ and $\Delta 12$ positions of the hydrocarbon chain. Oleic acid with only one double bond at the $\Delta 9$ position is more stable to oxidation than linoleic acid [2]. As a result, the oleic acid and linoleic acid (O/L) ratio is an important parameter for oil quality determination. By determining fatty acid composition the first two natural high-oleate peanut mutants (F435-2-1 and F435-2-2) were identified. The O/L ratio of the mutants reached nearly 40:1 [3]. Numerous high O/L peanut cultivars such as 'SunOleic 95R', 'SunOleic 97R', 'Florida-07' and some Chinese peanut varieties including Kainong176, and K17-15 were developed from F435 via traditional breeding efforts [4,5,6]. High oleic acid content in these mutants was controlled by two major recessive genes

[7]. Using Arabidopsis FAD2 cDNA as a probe, two homologous and non-allelic genes encoding $\Delta 12$ fatty acid desaturase were isolated and characterized in peanut [8,9]. $\Delta 12$ fatty acid desaturase is responsible for the conversion of oleic acid to linoleic acid in peanut [8,10]. One of these two genes was from A subgenome and designated as *AhFAD2A*. The mutant allele *fad2a* had G \rightarrow A transition at position 448 after the start codon. This caused a nonsynonymous amino acid substitution from aspartic acid to asparagine (D150N). This mutation resulted in a dysfunctional *AhFAD2A* desaturase. FAD2 gene from B subgenome was designated as *AhFAD2B*. The mutant allele *fad2b* had an "A" insertion at position 442 after the start codon. This insertion resulted in a frame shift and led to an inactive *AhFAD2B* desaturase [9,10]. There are only 11 base pair difference between *AhFAD2A* and *AhFAD2B*, resulting in a difference of 4 amino acids.

Based on the gene character of *AhFAD2A* and *AhFAD2B*, DNA markers were developed for marker-assisted selection (MAS) to enhance the efficiency of high oleic acid peanut breeding program. Cleaved amplified polymorphic sequence (CAPS) markers were first developed for *AhFAD2A* [11] and *AhFAD2B* detection [12]. Subsequently, allele-specific polymerase chain reaction (AS-PCR) method was developed [13] and further optimized [14]. PCR product sequencing

* Corresponding author.

E-mail address: xingjunw@hotmail.com (X. Wang).

Peer review under responsibility of Pontificia Universidad Católica de Valparaíso.

method was also used to distinguish *AhFAD2B* gene in the hybrids from crosses between normal cultivars and high oleate cultivars [15]. Real-time PCR amplification of *AhFAD2B* and *AhFAD2A* was used to discriminate wild type plants from the mutant alleles [16,17]. These various markers for *AhFAD2A* and *AhFAD2B* detection allow breeders and researchers to determine the genotypes of peanut lines with AABB, aabb and AaBb. There are nine genotypes in F_2 hybrids. It is difficult to distinguish all these genotypes within one experiment using the above-mentioned methods. Furthermore, utilization of restriction enzymes in CAPS and fluorescence-labeled probes in real-time PCR resulted in relatively high cost of these methods. Therefore, a simple, high throughput and low cost assay with high resolution is required.

KASP (Kompetitive Allele Specific PCR) is a novel competitive allele specific PCR for SNP genotyping assay based on dual FRET (Fluorescent Resonance Energy Transfer). In this method, the sample DNA is amplified with thermal cycler using allele specific primers. The allele specific primers are conjugated to fluorometric dye HEX and FAM at their 5' end, respectively. When FRET cassette primer is hybridized with DNA, fluorometric dye and quencher are separated, leading to emitting of the corresponding fluorescence. The genotype is easily detected by reading fluorescent signals. Owing to low cost, high throughput, and high specificity and sensitivity, KASP has been extensively used in massive SNP genotyping studies such as genome-wide SNP assay for rice genotyping [18], molecular markers-assisted breeding of wheat leaf rust resistance [19], and soybean cyst nematode resistance [20]. In peanut, KASP assay was firstly applied to screen on 94 genotypes using 96 SNPs, [21]. In this study, the KASP method was developed to detect all genotypes of *AhFAD2A* and *AhFAD2B* in different cross progenies.

2. Materials and methods

2.1. Plant materials

To develop peanut with high oleate and high yield for domestic consumption, cross was made between high oleate donors, Kainong176 (developed by Kaifeng Academy of Agriculture and Forestry), and normal oleate and high yield peanut cultivars, Huayu 31 (developed by Shandong Peanut Research Institute).

2.2. DNA extraction

About 0.1 g of a slice of cotyledonary tissue from parents, F_1 and F_2 mature seeds was obtained and placed with a 3.97 mm steel ball in a 1.5 mL tube. Tubes were immersed in liquid nitrogen and tissue was finely ground by intensely shaking the tube for 1 min. DNA extraction was performed according to the protocol of TIANGEN plant genomic DNA extraction kit (category No. DP-305).

2.3. KASP method

The KASP method was used to detect crossing parents, F_1 and BC_4F_2 seeds. The KASP reaction and its components are described at <http://www.lgcgenomics.com/genotyping/kasp-genotyping-reagents/how-does-kasp-work>. Sequences of allele-specific and common primers are listed in Table 1. 1.5 μ L $2\times$ KASP PCR mix, 35–45 ng of genomic DNA template, 0.16 μ M allele-specific forward primers, and 0.41 μ M reverse primers were incorporated into 3 μ L of KASP reaction. Amplification was performed in a Hydrocycler, water bath thermal cyclers, starting with 15 min at 94°C, a touchdown phase of 10 cycles at 94°C for 20 s and at 65°C for 60 s with a 1°C decrease in temperature per cycle, followed by 35 cycles of 94°C for 20 s and 55°C for 60 s. Once the thermal cycle is complete, BMG Omega F plate reader was used to read fluorescence signal. Fluorescence signal was acquired at 520 nm (green) and 556 nm (yellow) for 2 min at 25°C.

Table 1
Primers for KASP and CAPS.

Primer name	Sequence
<i>AhFAD2A</i> -Allele-1	5'-GTTTTGGGACAAACACTTCGTT-3'
<i>AhFAD2A</i> -Allele-2	5'-GTTTTGGGACAAACACTTCGTC-3'
<i>AhFAD2A</i> -common	5'-CGCCACCCTCCAACACC-3'
<i>AhFAD2B</i> -Allele-1	5'-CAAACACTTCGTCGGGTCT-3'
<i>AhFAD2B</i> -Allele-2	5'-CAAACACTTCGTCGGGTGCG-3'
<i>AhFAD2B</i> -common	5'-CCGCCACCCTCCAACACA-3'
Allele-1 Tail (FAM tail)	5'-GAAGGTGACCAAGTTCATGCT-3'
Allele-2 Tail (HEX tail)	5'-GAAGTTCGGAGTCAACGGATT-3'
aF19	5'-GATTACTGATTATTGACTT-3'
1056	5'-CCAACCCAAACCTTTCAGAG-3'
bF19	5'-CAGAACCATTAGCTTTG-3'
R1/FAD	5'-CTCTGACTATGCATCAG-3'

2.4. Validation of KASP using CAPS method

CAPS method [11,12] was used to validate the KASP results of *AhFAD2A* and *AhFAD2B*. Briefly, to detect *AhFAD2A* genotype, the genomic DNA was amplified with primers aF19 and 1056 (Table 1). Each reaction contained 1 μ L of DNA, 2 μ L of $10\times$ PCR buffer (Takara), 0.2 μ L of ExTaq DNA polymerase (Takara), 1.6 μ L of 2.5 mM dNTPs (Takara), 0.5 μ L of forward and reverse primers in 20 μ L reaction system. The PCR condition was 94°C for 5 min followed by 35 cycles of 94°C for 30 s, 51°C for 30 s, and 72°C for 1 min. The final extension step was at 72°C for 7 min. PCR products were digested with 1 U of *Hpy99I* restriction enzyme (New England Biolabs, Ipswich, MA) at 37°C for 4 h. The digested products were separated on a 2% (w/v) agarose gel and stained with ethidium bromide. To detect the *AhFAD2B* genotype, primer pairs bF19/R1(FAD) (Table 1) amplified the coding region of *AhFAD2B* genes using a total PCR reaction volume of 20 μ L containing 0.5 μ L DNA, 0.5 μ L FastPfu DNA polymerase (Transgene Biotech), 4.0 μ L of $5\times$ Pfu buffer, 0.4 μ L of 10 mM dNTPs (Transgene Biotech), and 0.5 μ L forward and reverse primers R1(FAD), respectively. Amplification conditions were initiated at 94°C for 5 min for denaturation; 35 cycles of 94°C for 45 s, 52°C for 45 s, 72°C for 60 s; final extension at 72°C for 7 min. Eight microliters of the amplification products was digested by 0.2 μ L or 2 U of *Hpy188I* (New England Biolabs, Ipswich, MA, USA). The digestion was performed at 37°C overnight. Digested products were separated on a 2% (w/v) agarose gel.

3. Results and discussion

3.1. Genotyping *AhFAD2A* and *AhFAD2B* by KASP

The genotypes of *AhFAD2A* and *AhFAD2B* genes in normal oleic peanut and high oleate peanut were designated as AABB and aabb, respectively. To develop the KASP method allele-specific primers were designed based on the sequences of wild type and mutant *AhFAD2A* and *AhFAD2B* genes (Table 1). The KASP assay mix contains three assay-specific non-labeled oligos: two allele-specific forward primers and one common reverse primer. The allele-specific forward primers differentiated the mutant allele from wild type allele. Due to the high sequence identity of coding region of *AhFAD2A* and *AhFAD2B* genes, the primers are the key to successfully distinguish these two genes. Fortunately, there is a SNP between these two genes at position 432 from the 5' end of the open reading frame (C in *FAD2A* and A in *FAD2B*) [10]. Therefore, according to the competitive allele specific PCR theory of KASP, 3' end of the common reverse primer was designed at this SNP position to discriminate A and B subgenomes (Fig. 1).

The allele-specific primers each harbored a unique tail sequence that corresponds with a universal FRET (fluorescence resonant energy transfer) cassette; *Ahfad2a/Ahfad2b*-Allele-1 Tail primer, the mutant




Fig. 1. Locations of genetic mutations and primer design in both *AhFAD2A* and *AhFAD2B* alleles. Different primer sites were shown with different colors. All primer sequences were shown in Table 1.

allele-specific primer, was labeled with FAM™ dye and *AhFAD2A/AhFAD2B*-Allele-2 Tail primer, the WT allele-specific primer, was labeled with HEX™ dye. Bi-allelic discrimination is achieved through the competitive binding of these two allele-specific forward primers. If the genotype at a given SNP is aa or bb homozygous, only FAM fluorescent signal could be generated. If the genotype at a given SNP is AA or BB homozygous, only HEX fluorescent signal could be obtained. If the genotype is heterozygous, a mixed fluorescent signal was produced. KASP method was used to genotype the seeds from the male parent (aabb), female parent (AABB) and F₁ (AaBb) in this study. As shown in Fig. 2, three different genotypes were clearly separated, and the dots of each genotype were closely assembled. Therefore, our results demonstrated that the developed KASP could detect the genotypes of peanut *AhFAD2A* and *AhFAD2B* genes.

3.2. Validation of KASP using CAPS

CAPS assay was successfully developed to detect the genotype of *AhFAD2A* and *AhFAD2B* [11,12]. In this study, we confirmed some of the KASP results by CAPS analysis. The recognition site sequence of *Hpy99I* is 5'-CGWCG-3' (W = A/T). This restriction site was found in the ORF of the wild-type *FAD2A* but not in high O/L mutant peanut lines [22,23]. Therefore, the 826 bp PCR product amplified from *AhFAD2A* gene of WT peanut could be recognized by *Hpy99I* and cut into two fragments, 598 bp and 228 bp in length (Fig. 3a). In the *Ahfad2a* mutant lines, the PCR product could not be recognized by *Hpy99I* and then no digestion product could be observed. Incomplete digestion could be observed in WT peanut in part due to the low fidelity of the Taq polymerase used for PCR amplification [11]. The

recognition site sequence of *Hpy188I* is 5'-TCNGA-3'. Due to the insertion of an "A" between 441_442 resulted in the "TCAGA" in the *Ahfad2b* mutant allele. The mutant allele, therefore, can be digested by *Hpy188I* at this position, whereas the *AhFAD2B* wild-type and *AhFAD2A* alleles cannot be digested (Fig. 3b). DNA was extracted from the male parent, the female parent, and 20 F₁ seeds, which have been analyzed by KASP. After PCR amplification and digestion with *Hpy99I* and *Hpy188I*, genotypes of 22 samples were all consistent with the results obtained from KASP. These results together indicated that we successfully developed KASP assay method to efficiently and accurately distinguish *AhFAD2A* and *AhFAD2B* of the WT, mutant and F₁ hybrids.

3.3. Application of KASP in genotyping of BC₄F₂ generation

Due to the nature of allotetraploid, the cross between homozygote high oleic acid peanut (aabb) and normal peanut cultivar (AABB) produces 9 genotypes in F₂ or BC₄F₂ generations. These genotypes are AABB, AaBB, AABb, AaBb, aaBB, aaBb, AAbb, Aabb and aabb, respectively. Furthermore, the number of F₂ or BC₄F₂ seeds is about 30 times of F₁ or BC₄F₁ (If one F₁ plant could produce 30 seeds). Compared to the other detection method, KASP is simple, low cost and efficient. In this study, KASP was successfully used to examine the genotypes of BC₄F₂ segregating populations. Out of 179 BC₄F₂ seeds detected, the genotype of 18 F₂ seeds was aabb, and the genotype of 25 seeds was AaBb (Supplementary Table 1 and Supplementary Table 2). Therefore, we successfully developed KASP assay to efficiently and accurately distinguish all forms of *AhFAD2A* and *AhFAD2B* in peanut.




Fig. 2. Genotyping results of *AhFAD2A* (a) and *AhFAD2B* (b) by KASP. The scatter plot with axes x and y represents allelic discrimination of *FAD2A* or *FAD2B* genotypes. The red, green and blue dots represent the mutant homozygous, heterozygous and wild-type homozygous, respectively.




Fig. 3. Genotyping results of *AhFAD2A* and *AhFAD2B* by CAPS. (a) *Hpy99I* digestion of PCR products; (b) *Hpy188I* digestion of PCR products. M referred to DL5000 DNA Marker; Lane 1 and Lane 2 showed the genotypes of Huayu31 and Kainong176, respectively; Lanes 3–22 showed the genotypes of 20 F_1 seeds, respectively.

4. Conclusions

In this research, KASP has been successfully developed to detect the genotype of *AhFAD2A* and *AhFAD2B* in peanut. Due to high accuracy, saving time and resources, this method will greatly facilitate breeding progress of high oleate peanut. It is more suitable for determining *AhFAD2* genotype than other methods.

Conflict of interests

The authors declare that they have no competing interests.

Financial support

This study was financially supported by MARS Company, Shandong Province Germplasm Innovation and Utilization Project, the National High Technology Research and Development Program of China (2013AA102602), the National Natural Science Foundation of China (31471526), Shandong Provincial Natural Science Foundation (BS2013SW006), and the Young Talents Training Program of Shandong Academy of Agricultural Sciences.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <http://dx.doi.org/10.1016/j.ejbt.2016.10.010>.

References

- Ahmed E, Young C. Composition, quality, and flavor of peanuts. In: Pattee HE, Young CT, editors. Peanut science and technology. Yoakum: American Peanut Research and Education Society. Inc; 1982. p. 655–88.
- Schwartzbeck JL, Jung S, Abbott AG, Mosley E, Lewis S, Pries GL, et al. Endoplasmic oleoyl-PC desaturase references the second double bond. *Phytochemistry* 2001;57: 643–52. [http://dx.doi.org/10.1016/S0031-9422\(01\)00081-4](http://dx.doi.org/10.1016/S0031-9422(01)00081-4).
- Norden A, Gorbet D, Knauff D, Young C. Variability in oil quality among peanut genotypes in the Florida breeding program 1. *Peanut Sci* 1987;14:7–11. <http://dx.doi.org/10.3146/i0095-3679-14-1-3>.
- Gorbet DW, Knauff D. Registration of 'SunOleic 95R' peanut. *Crop Sci* 1997;37:1392.
- Gorbet DW, Knauff D. Registration of 'SunOleic 97R' Peanut. *Crop Sci* 2000;40: 1189–90. <http://dx.doi.org/10.2135/cropsci2000.0032rcv>.
- Gorbet DW, Tillman BL. Registration of 'Florida-07' peanut. *J Plant Regist* 2009;3: 14–8. <http://dx.doi.org/10.3198/jpr2008.05.0276rc>.
- Moore K, Knauff D. The inheritance of high oleic acid in peanut. *J Hered* 1989;80: 252–3.
- Jung S, Powell G, Moore K, Abbott A. The high oleate trait in the cultivated peanut (*Arachis hypogaea* L.). II. Molecular basis and genetics of the trait. *Mol Gen Genet* 2000;263:806–11. <http://dx.doi.org/10.1007/s004380000243>.
- Lopez Y, Nadaf HL, Smith OD, Connell JP, Reddy AS, Fritz AK. Isolation and characterization of the $\Delta 12$ -fatty acid desaturase in peanut (*Arachis hypogaea* L.) and search for polymorphisms for the high oleate trait in Spanish market-type lines. *Theor Appl Genet* 2000;101:1131–8. <http://dx.doi.org/10.1007/s001220051589>.
- Jung S, Swift D, Sengoku E, Patel M, Teule F, Powell G, et al. The high oleate trait in the cultivated peanut (*Arachis hypogaea* L.). I. Isolation and characterization of two genes encoding microsomal oleoyl-PC desaturases. *Mol Gen Genet* 2000;263: 796–805. <http://dx.doi.org/10.1007/s004380000244>.
- Chu Y, Ramos L, Holbrook C, Ozias-Akins P. Frequency of a loss-of-function mutation in Oleoyl-PC Desaturase (*AhFAD2A*) in the mini-core of the US peanut germplasm collection. *Crop Sci* 2007;47:2372–8. <http://dx.doi.org/10.2135/cropsci2007.02.0117>.
- Chu Y, Holbrook CC, Ozias-Akins P. Two alleles of *ahFAD2B* control the high oleic acid trait in cultivated peanut. *Crop Sci* 2009;49:2029–36. <http://dx.doi.org/10.2135/cropsci2009.01.0021>.
- Chen Z, Wang ML, Barkley NA, Pittman RN. A simple allele-specific PCR assay for detecting *FAD2* alleles in both A and B genomes of the cultivated peanut for high-oleate trait selection. *Plant Mol Biol Rep* 2010;28:542–8. <http://dx.doi.org/10.1007/s11105-010-0181-5>.
- Yu HT, Yang WQ, Tang YY, Wang XZ, Wu Q, Hu DQ, et al. An AS-PCR assay for accurate genotyping of *FAD2A/FAD2B* genes in peanuts (*Arachis hypogaea* L.). *Grasas Aceites* 2013;64:395–9. <http://dx.doi.org/10.3989/gya.118712>.
- Wang CT, Yu SL, Zhang SW, Wang XZ, Tang YY, Zhang JC, et al. Novel protocol to identify true hybrids in normal oleate \times high oleate crosses in peanut. *Electron J Biotechnol* 2010;13. <http://dx.doi.org/10.2225/vol13-issue5-fulltext-18>.
- Barkley NA, Chamberlin KDC, Wang ML, Pittman RN. Development of a real-time PCR genotyping assay to identify high oleic acid peanuts (*Arachis hypogaea* L.). *Mol Breed* 2010;25:541–8. <http://dx.doi.org/10.1007/s11032-009-9338-z>.
- Barkley NA, Wang ML, Pittman RN. A real-time PCR genotyping assay to detect *FAD2A* SNPs in peanuts (*Arachis hypogaea* L.). *Electron J Biotechnol* 2011;14(1): 9–10.
- McCouch SR, Zhao K, Wright M, Tung CW, Ebana K, Thomson M, et al. Development of genome-wide SNP assays for rice. *Breed Sci* 2010;60:524–35. <http://dx.doi.org/10.1270/jsbbs.60.524>.
- Neelam K, Brown-Guedira G, Huang L. Development and validation of a breeder-friendly KASPar marker for wheat leaf rust resistance locus *Lr21*. *Mol Breed* 2013; 31:233–7. <http://dx.doi.org/10.1007/s11032-012-9773-0>.
- Shi Z, Liu S, Noe J, Arelli P, Meksem K, Li Z. SNP identification and marker assay development for high-throughput selection of soybean cyst nematode resistance. *BMC Genomics* 2015;16:314. <http://dx.doi.org/10.1186/s12864-015-1531-3>.
- Khera P, Upadhyaya HD, Pandey MK, Roorkiwal M, Sriswathi M, Janila P, et al. Single nucleotide polymorphism-based genetic diversity in the reference set of peanut (*Arachis* spp.) by developing and applying cost-effective competitive allele specific polymerase chain reaction genotyping assays. *Plant Genome* 2013;6. <http://dx.doi.org/10.3835/plantgenome2013.06.0019>.
- Bruner AC, Jung S, Abbott AG, Powell GL. The naturally occurring high oleate oil character in some peanut varieties results from reduced oleoyl-PC desaturase activity from mutation of aspartate 150 to asparagine. *Crop Sci* 2001;41:522–6. <http://dx.doi.org/10.2135/cropsci2001.412522x>.
- Patel M, Jung S, Moore K, Powell G, Ainsworth C, Abbott A. High-oleate peanut mutants result from a MITE insertion into the *FAD2* gene. *Theor Appl Genet* 2004; 108:1492–502. <http://dx.doi.org/10.1007/s00122-004-1590-3>.